



Review

Trabecular bone score (TBS) as a new complementary approach for osteoporosis evaluation in clinical practice[☆]



N.C. Harvey^{a,b,*}, C.C. Glüer^c, N. Binkley^d, E.V. McCloskey^e, M.-L. Brandi^f, C. Cooper^{a,b,g}, D. Kendler^h, O. Lamyⁱ, A. Laslop^j, B.M. Camargos^k, J.-Y. Reginster^l, R. Rizzoli^m, J.A. Kanis^{e,**}

^a MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton, UK

^b NIHR Southampton Nutrition Biomedical Research Centre, University of Southampton and University Hospital Southampton NHS Foundation Trust, Southampton, UK

^c Sektion Biomedizinische Bildgebung, Klinik für Radiologie und Neuroradiologie, Universitätsklinikum Schleswig-Holstein, Campus Kiel, Kiel, Germany

^d Osteoporosis Clinical Research Program, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

^e Centre for Metabolic Bone Diseases, University of Sheffield, Sheffield, UK

^f Department of Surgery and Translational Medicine, University of Florence, Florence, Italy

^g NIHR Musculoskeletal Biomedical Research Unit, University of Oxford, Oxford, UK

^h Department of Medicine, University of British Columbia, Vancouver, Canada

ⁱ Bone Unit, University Hospital, Lausanne, Switzerland

^j Scientific Office, Austrian Agency for Health and Food Safety, Vienna, Austria

^k Unidade de Densitometria Óssea, Densimater Rede Materdei de Saúde, Belo Horizonte, MG, Brazil

^l Department of Public Health, Epidemiology and Health Economics, University of Liege, Liege, Belgium

^m Service of Bone Diseases, Geneva University Hospitals and Faculty of Medicine, Geneva, Switzerland

ARTICLE INFO

Article history:

Received 21 April 2015

Accepted 12 May 2015

Available online 16 May 2015

Edited by R. Baron

Keywords:

Osteoporosis

Epidemiology

Trabecular bone score (TBS)

Fragility fracture

Bone mineral density

FRAX

ABSTRACT

Trabecular bone score (TBS) is a recently-developed analytical tool that performs novel grey-level texture measurements on lumbar spine dual X-ray absorptiometry (DXA) images, and thereby captures information relating to trabecular microarchitecture. In order for TBS to usefully add to bone mineral density (BMD) and clinical risk factors in osteoporosis risk stratification, it must be independently associated with fracture risk, readily obtainable, and ideally, present a risk which is amenable to osteoporosis treatment. This paper summarizes a review of the scientific literature performed by a Working Group of the European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis. Low TBS is consistently associated with an increase in both prevalent and incident fractures that is partly independent of both clinical risk factors and areal BMD (aBMD) at the lumbar spine and proximal femur. More recently, TBS has been shown to have predictive value for fracture independent of fracture probabilities using the FRAX® algorithm. Although TBS changes with osteoporosis treatment, the magnitude is less than that of aBMD of the spine, and it is not clear how change in TBS relates to fracture risk reduction. TBS may also have a role in the assessment of fracture risk in some causes of secondary osteoporosis (e.g., diabetes, hyperparathyroidism and glucocorticoid-induced osteoporosis). In conclusion, there is a role for TBS in fracture risk assessment in combination with both aBMD and FRAX.

© 2015 Elsevier Inc. All rights reserved.

Contents

Introduction	217
Osteoporosis, fragility fractures and risk assessment	217
Trabecular bone score: definition	217
Search strategy	217
Does TBS predict osteoporosis-related fracture risk?	217
Cross-sectional studies	218
Prospective studies	218

[☆] A consensus report of a European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO) Working Group.

* Correspondence to: N.C. Harvey, MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton, UK.

** Corresponding author.

E-mail addresses: nch@mrc.soton.ac.uk (N.C. Harvey), w.j.pontefract@sheffield.ac.uk (J.A. Kanis).

Is TBS a potential adjunct to FRAX® probability?	220
Is TBS responsive to treatment?	220
Does TBS have a role in secondary osteoporosis?	222
Clinical and subclinical hypercortisolism	222
Type-2 diabetes	222
Primary hyperparathyroidism	222
Osteoarthritis	222
Conclusions	222
Disclosures	223
Acknowledgments	223
References	223

Introduction

Osteoporosis, fragility fractures and risk assessment

Measurements of bone mineral density (BMD) are a central component of any provision that arises from the definition of osteoporosis, agreed internationally as: a progressive systemic skeletal disease characterized by low bone mass and microarchitectural deterioration of bone tissue, with a consequent increase in bone fragility and susceptibility to fracture [1]. This definition captures the notion that low areal BMD (aBMD) is an important component of fracture risk, but that other bone abnormalities contribute to skeletal fragility. The conceptual description of osteoporosis thus centres both on the assessment of bone mass and quality, specifically bone microstructure. Until recently, there were no satisfactory clinical means to assess bone microstructure non-invasively, so that the operational diagnosis of osteoporosis is based on the measurement of aBMD. Osteoporosis is so-defined as a femoral neck aBMD 2.5 SD or more below the young adult female mean (T-score ≤ -2.5) [2,3]. The same T-score derived at other sites is widely used in clinical practice (e.g., lumbar spine, total hip, distal radius).

A consequence of this operational definition, which identifies the small proportion of the population at highest risk, is that the greater number of individuals above this threshold, although individually at lower risk, contributes the greater number of fractures to the total burden. Indeed, the majority of fragility fractures occur in patients who have an aBMD T-score > -2.5 . In other words, the detection rate for these fractures (sensitivity) is low [4], which is why widespread population-based screening is not generally recommended in women at menopause [2,5]. Thus, factors other than bone mass influence bone strength and fracture risk, including microarchitectural deterioration of bone tissue, as given in the conceptual definition of osteoporosis. Additional skeletal and extra-skeletal factors, such as bone geometry, micro-damage, mineralization, bone turnover, age, and a large range of clinical risk factors, including family history, prior fracture and fall risk, contribute to the overall assessment of fracture risk [6–9]. Several of these additional factors are captured by FRAX®. FRAX estimates the 10-year probability of hip and major osteoporotic fracture based on the individual's risk factor profile [4]. Apart from BMD, FRAX does not capture other skeletal determinants of bone strength that improve upon or are at least partly independent of aBMD [10]. Several such determinants are the subject of clinical research [11–18] using novel imaging techniques, such as quantitative computed tomography (QCT) and high resolution (peripheral) QCT [19,20], and minimally invasive approaches for probing bone material properties, notably microindentation techniques [21]. Although there is evidence of their predictive ability for fracture [22,23], none of these modalities appears to reliably outperform aBMD in the prediction of the various types of osteoporotic fractures, and their general lack of availability and validation in the clinical setting means that an adjunctive role alongside DXA-measured aBMD is unlikely to be feasible in most settings in the near future. In contrast, trabecular bone score (TBS) is a novel imaging technique,

based on standard DXA images, and appears to constitute an index of bone texture that provides skeletal information additional to the standard aBMD results [24].

Trabecular bone score: definition

TBS has emerged as a novel grey-level texture measurement that uses experimental variograms of 2D projection images, quantifying variation in grey-level texture from 1 pixel to the adjacent pixels. TBS is not a direct measurement of bone microarchitecture but it is related to 3D bone characteristics such as the trabecular number, the trabecular separation and the connectivity density [25,26]. An elevated TBS appears to represent strong, fracture-resistant microarchitecture, whilst a low TBS reflects weak, fracture-prone microarchitecture. As such, there is evidence that TBS can differentiate between two 3-dimensional (3D) microarchitectures that exhibit the same bone density, but different trabecular characteristics. TBS is generally obtained by re-analysis of AP lumbar spine DXA images, which allows direct comparison with aBMD and application to existing datasets. This latter opportunity has led to a rapid rise in published research assessing its potential role in the assessment and management of osteoporosis.

Lumbar TBS, like aBMD, is an age dependent variable. Little change in TBS is observed between the ages of 30 and 45 years. Thereafter, a progressive decrease is observed with advancing age [27], which is more marked in women than in men. The percentage decrease with age is similar to that for lumbar spine aBMD, as is the short term reproducibility [25].

This paper reports the findings of a European Society for Clinical and Economic Aspects of Osteoporosis and Osteoarthritis (ESCEO) Working Group, which first convened in September 2014 with the aim of comprehensively assessing the evidence supporting the use of TBS in clinical practice. More specifically, this report reviews the potential value of TBS as an independent adjunct to risk assessment using DXA aBMD and/or FRAX in settings such as post-menopausal and secondary osteoporosis, and its potential use in assessment of response to treatment.

Search strategy

A Medline search for publications with the terms trabecular bone score or TBS was undertaken in September 2014. Published articles in English and French were extracted. Papers in abstract form were not included except where the authors supplied a full copy of the submitted manuscript. A total of 479 papers were identified of which 67 manuscripts were considered relevant and the full publication reviewed. The search was subsequently updated in February 2015 and a total of 73 papers were reviewed.

Does TBS predict osteoporosis-related fracture risk?

To date, eighteen studies assessing fracture risk in post-menopausal women have been published. Of these, eleven were cross-sectional [28–38] and seven prospective, including a meta-analysis [25,39–44].

Cross-sectional studies

The eleven cross-sectional studies are summarized in Table 1. The studies all were published from 2009 onward to January 2015. Sample sizes ranged from 135 to 3069, with the majority between 150 and 300 subjects. Six of the studies exclusively involved post-menopausal women, whilst one included only men over 40 years [31], and another both men and women over 50 years who had suffered at least one prior fracture [33]. There was variation in which fractures were assessed and how the fractures were ascertained. For example, five studies used self-report for all fractures, though vertebral fracture assessments (VFA) were used to identify vertebral fractures in three studies. In every study, however, and for every fracture type, there were significantly increased odds of prior fracture among those with low TBS even when adjusted for age, lumbar aBMD, body mass index, and clinical risk factors. Moreover, in one study which reported area (AUC) under receiver operating curves (ROC) as a combined measure of sensitivity and specificity, the AUC for TBS was greater than for aBMD of the spine (0.67 vs. 0.54, $p = 0.035$) but not for hip aBMD for subjects in the non-osteoporotic aBMD range [33]. However, in a further study which reported AUC values, these were similar for total hip BMD and total hip BMD with TBS (0.80 versus 0.81 respectively) [36].

Prospective studies

Cross-sectional studies may not accurately quantify the utility of a measure in the longitudinal prediction of incident fractures. However, data from prospective studies have supported the findings from the

cross-sectional investigations. Of the six prospective single studies, one was conducted in France [28], one in the UK, France and Germany [40], three in Canada [25,43,45,54], and one in Japan [41] (Table 2). These cohorts tended to be larger than those in the cross-sectional studies, with the smallest with 560 subjects and the largest over 33,000 post-menopausal women; follow-up ranged from under 5 to over 8 years.

By far the largest published study assessing TBS to date was conducted in the Canadian province of Manitoba [46,47], comprising 29,407 postmenopausal women (all over 50 years, mean 65.4 years) living in and around the capital city of Winnipeg [25]. At five years of follow-up, there were 1668 incident major osteoporotic fractures, including 439 vertebral fractures and 293 hip fractures. Lumbar spine aBMD and TBS predicted fractures equally well, and the combination of both performed better than either individually. Interestingly, although aBMD-lumbar spine and aBMD-total hip were closely correlated ($r = 0.72$), aBMD-lumbar spine and TBS-lumbar spine were only weakly correlated ($r = 0.32$), with only 10% of the variance in one explained by variance in the other. Similarly weak to moderate correlations between aBMD and TBS have been reported by several others [48–53], and may relate to measurement of different bone properties, or measurement variability. When various models adjusted for age and clinical risk factors were tested for overall accuracy using receiver operating curves (ROC) and AUC estimates, the model combining aBMD-total hip and TBS-spine performed best, with AUC values for clinical vertebral and hip fractures of 0.73 (95% CI: 0.71, 0.75) and 0.82 (95% CI: 0.79, 0.84), respectively. For femoral neck, total hip and lumbar aBMD, the AUC improved

Table 1
Cross-sectional case control studies that examine TBS and fracture risk.

First author	Year published	Location	Subjects	Outcome(s)	TBS		BMD		Adjustments
					OR/SD decrease	95% CI	OR/SD decrease	95% CI	
Pothuau [34]	2009	Switzerland	135 PM females	Vertebral Fx	2.66	1.46, 4.85	NR	NR	None
			France	45 cases with Fx	hip Fx	NR	NR	NR	NR
Winzenrieth [38]	2010	France	90 BMD-matched controls	MOF	1.95	1.31, 2.89	NR	NR	None
			243 PM females	Vertebral Fx	1.97	1.31, 2.96	1.63	1.20, 2.22	Weight
Rabier [35]	2010	France	81 cases with vertebral Fx	vertebral Fx	3.81	2.17, 6.72	2.48	1.61, 3.83	Weight
			162 age-matched controls						
Lamy [30]	2012	Switzerland	168 PM females	All Fx (self-report)	1.4	1.1, 1.7	1.3	1.1, 1.6	Age, BMI
			42 cases vertebral Fx						
Del Rio [28]	2013	Spain	126 age-matched controls	vertebral Fx (VFA)	2.0	1.4, 3.0	1.8	1.2, 2.5	age, BMI
			631 PM women	MOF	1.9	1.4, 2.5	1.6	1.2, 2.1	age, BMI
Vasic [37]	2013	Serbia	164 with Fx (53 VF, 107 MOF)	Femoral neck Fx (self-report)	1.71	1.15, 2.79	1.94	1.35, 2.79	Age
			467 controls						
Leib [32]	2014	USA	191 PM women	All Fx (self-report)	1.28	1.13, 1.46	NR	NR	Age, LS BMD, weight, smoking, maternal history of hip fracture
			83 cases with Fx						
Krueger [29]	2014	USA	108 non-matched controls	fracture (self-report)	2.46	1.9, 3.1	1.36	1.2, 1.6	Age, BMI
			1031 women						
Leib [31]	2014	USA	271 with Fx	low-energy Fx (self-report)	1.55	1.09, 2.20	NR	NR	Age, LS BMD
			2165 women						
Nassar [33]	2014	France	289 cases with Fx	vertebral Fx (VFA) vs. non-vertebral Fx	NR	NR	NR	NR	NR
			1876 non-matched controls						
Touvier [36]	2015	France	271 age-matched controls	AUC TBS > AUC BMD (0.67 vs. 0.54, $p = 0.04$)	2.25	1.58, 3.29	1.70	1.18, 2.54	Age
			180 men						
			135 age/LS-BMD matched controls	Fragility fracture (orthopaedic clinic)	2.25	1.58, 3.29	1.70	1.18, 2.54	Age
			280 women, 82 men > age 50						
			133 vertebral Fx, 186 hip Fx						
			176 non-vertebral Fx, non-hip Fx						
			255 women, 40–92 years						
			79 fragility fracture						
			176 controls						

PM = postmenopausal, Fx = fracture; MOF = major osteoporotic fracture; NR = not recorded.

Table 2
Prospective cohort studies that have examined lumbar TBS and fracture risk.

First author, ref number	Year published	Location	Subjects	Outcome	Follow-up [years]	Lumbar spine TBS HR or RR/SD ^a	95% CI	Lumbar spine BMD HR or RR/SD ^a	95% CI	Total hip BMD HR or RR/SD ^a	95% CI	Adjustments
Hans [25]	2011	Canada	29,407 PM women 5.7% OP Fx (1.5% clinical VFx); 1.0% hip Fx	Clinical VF hip Fx MOF	4.7	1.45 1.46 1.35	1.32, 1.58 1.30, 1.63 1.29, 1.42	1.72 1.31 1.47	1.55, 1.91 1.16, 1.48 1.39, 1.55	1.75 2.55 1.67	1.58, 1.96 2.22, 2.93 1.58, 1.76	Age Age Age
Boutroy [39]	2013	France	OLEFY cohort 560 PM women 94 with Fx at any site	Clinical VF radiographic VF fragility Fx	8	1.34	1.04, 1.73	1.3	1.06, 1.58	1.99	1.52, 2.62	Age, weight
Briot [40]	2013	France UK Germany	OPUS cohort 1007 PM women 8.1% with OP Fx; 4.6% with VF	Radiographic VF clinical OP Fx	6	1.54 1.62	1.17, 2.03 1.30, 2.01	1.75 1.47	1.25, 2.48 1.16, 1.89	1.73 1.65	1.26, 2.38 1.30, 2.11	None
Iki [41]	2014	Japan	665 PM women 92 with VF	VF (VFA)	8.3	1.54 OR per SD decrease	1.17, 2.02	1.27 OR per SD decrease	1.02, 1.59	NR	NR	Age, LS BMD
Leslie [54]	2014a	Canada	3620 men > 50 (mean age 67.6) 183 MOF (91 VF, 46 hip Fx)	MOF hip Fx VF	4.5	1.22 1.6 1.12	1.60, 1.41 1.21, 2.11 0.91, 1.38	1.14 1.29 1.26	1.02, 1.28 1.03, 1.60 1.09, 1.47	1.66 2.84 1.2	1.40, 1.98 1.95, 4.12 0.95, 1.51	FRAX score, treatment
Leslie [43]	2014b	Canada	33,352 women 40–100 years	MOF Death		1.36 1.32	1.30, 1.42 1.26, 1.39	NR	NR	NR		Time since baseline and age
McCloskey [42]	2015	Multinational	17,809 men & women 298 with hip 1109 with MOF	MOF hip Fx Men MOF Men hip Fx Women MOF Women hip Fx	6.7	1.32 1.28 1.35 1.27 1.31 1.29	1.24, 1.41 1.13, 1.45 1.21, 1.49 1.06, 1.53 1.21, 1.42 1.09, 1.52	NR	NR	NR		FRAX score, time since baseline and age

PM = postmenopausal, Fx = fracture; MOF = major osteoporotic fracture; VFx = vertebral fracture; NR = not recorded.

^a Hazard ratio or relative risk per SD decrease in predictor unless otherwise stated.

significantly ($p < 0.001$) with the inclusion of TBS. Fracture rate increased from highest to lowest third of TBS for subjects across the range of aBMD.

A smaller longitudinal study, conducted in France, was the OFELY (Osteoporose dans les Femmes de Lyon) study involving 560 postmenopausal women of mean age 65.3 years recruited in 2000 and 2001 and followed for a mean 7.8 years [39]. In this cohort there were 112 incident fragility fractures that occurred in 92 women. Adjusted for age and prior fracture, TBS, lumbar spine and total hip aBMD were predictive of fracture occurrence. As with the Manitoba study, the AUC for a combined model of lumbar aBMD and TBS was better than that for either test used alone. TBS appeared to have the discriminative ability only in those women with normal or osteopenic aBMD but not for women in the osteoporotic aBMD range. For example, the incidence of fracture in osteopenic women with a TBS value in the lowest quartile was 25% compared to 13% in osteopenic women with TBS values in the remaining quartiles.

The Japanese Population-based Osteoporosis Cohort Study (JPOS) prospectively followed 665 Japanese women over the age of 50 years (mean age 64.1 years) for evidence of new morphometric vertebral fractures (VF), over approximately ten years [41]. During follow-up, 92 new vertebral fractures were documented by vertebral fracture assessment (VFA). In contrast to the OFELY study, there was an increased rate of VF from highest to lowest third of TBS score irrespective of BMD; i.e., also in those women in the osteoporotic aBMD range; the combination of aBMD and TBS increased overall performance of VF risk estimation over either used alone.

In the Osteoporosis and Ultrasound Study (OPUS), 1007 postmenopausal women (mean age 65.9 years) were recruited from three European centres, one each in France, Germany and the UK, and followed for an average of 6.0 years [40]. Over those six years, there were 82 incident clinical fractures and 46 incident vertebral fractures identified by radiography. TBS and aBMD at lumbar spine, femoral neck or total hip were similarly predictive of fragility fractures, but the combination of TBS and aBMD performed better than either measure alone, both in osteoporotic and non-osteoporotic women.

Only one longitudinal study on TBS has data for men (Table 2), with 3620 men over age 50 years (mean age 67.6 years) in the Manitoba cohort, followed for an average of 4.5 years and sustaining 183 major osteoporotic fractures [54]. TBS was predictive of hip, vertebral or major fractures. However, predictive power remained significant only for hip fracture when adjusted for FRAX score, osteoporosis treatment, and aBMD. There was some evidence of an interaction between aBMD and TBS such that the relationship between TBS and incident fracture appeared stronger in those with an aBMD T-score < -2.5 .

A recent meta-analysis of 14 prospective population-based cohorts comprising 17,809 men and women (59% women) examined fracture risk expressed as a gradient of risk (GR, the increase in fracture risk/SD difference in TBS). The GR of TBS for hip fracture or other major osteoporotic fracture (clinical spine, distal forearm or proximal humerus fracture) ranged from 1.31 to 1.54 depending on age and fracture outcome with no difference between men and women [42].

These data support the utility of low lumbar TBS as a determinant of fracture risk, at least partly independent of aBMD. The consistency of the predictive value for fracture in international cohorts argues for its clinical use with aBMD.

Is TBS a potential adjunct to FRAX® probability?

FRAX is widely used as a fracture risk assessment tool and the question arises whether TBS might serve as an adjunct to FRAX risk factors in the stratification of fracture risk. For TBS to be considered clinically useful as a FRAX modifier, it should be at least partly independent of lumbar spine aBMD, femoral neck aBMD and FRAX clinical risk factors (CRFs). An assessment of the relationship between CRFs, aBMD and TBS was undertaken in the Manitoba cohort, in which 33,352 women aged

40–100 years (mean 63 years) with baseline DXA measurements of lumbar spine TBS and femoral neck aBMD were studied [43]. Over a mean of 4.7 years follow-up, 1754 women died and 1872 sustained one or more major osteoporotic fracture. Lower TBS predicted increased risk of incident fracture (HR/SD decrease: 1.36; 95% CI: 1.30, 1.42, $p < 0.001$) and a 32% increase in death rate (HR/SD decrease: 1.32; 95% CI: 1.26, 1.39, $p < 0.001$). Lumbar spine TBS remained a significant predictor of major osteoporotic fracture (HR/SD decrease: 1.18; 95% CI: 1.12, 1.23) after adjustment for significant clinical risk factors and femoral neck aBMD. Low TBS (10th percentile) increased fracture risk 1.5 to 1.6-fold relative to high TBS (90th percentile), after accounting for competing mortality, across a wide range of ages and femoral neck T-scores.

Similar results were found in 3620 men from the Manitoba cohort at the age of 50 years or more (mean 67.6 years) [54]. Over a mean follow-up of 4.5 years, 183 men sustained major osteoporotic fractures and 91 clinical vertebral fractures. Lower lumbar spine TBS was observed in fracture versus non-fracture men for all fracture categories, and the AUC for incident fracture discrimination with TBS was significantly better than by chance (major osteoporotic fracture AUC = 0.59, $p < 0.001$; hip fracture AUC = 0.67, $p < 0.001$; clinical vertebral fracture AUC = 0.57, $p = 0.032$). After adjustment for FRAX without aBMD and osteoporosis treatment, TBS predicted major osteoporotic fracture and hip fracture (but not clinical vertebral fracture), and remained a predictor of hip fracture (but not major osteoporotic fracture) after further adjustment for hip or spine aBMD.

More recently, Manitoba data have been used to derive an adjustment factor to alter FRAX probabilities when accounting for TBS [55]. Data from 33,352 women between the ages of 40 and 100 years in Manitoba were used, including baseline DXA measures of femoral neck aBMD and lumbar spine TBS. Hazard functions for risk of osteoporotic fracture without hip fracture, hip fracture, and death (since risk of death competes with risk of fracture) were used to compute the adjustment factor. Femoral neck aBMD, clinical risk factors and TBS all contributed independent predictive value to the models, and TBS modified the 10-year probability of fracture outcomes generated from clinical risk factors and aBMD in FRAX-like models. Thus, for example, in an 80-year old woman, with a femoral neck T score of -2 , and BMI 27 kg/m², the 10-year probability of major osteoporotic fracture was 16.5%. If her TBS was found to be low (10th percentile), this would increase her fracture probability to 18.0%; conversely if TBS was found to be high (90th percentile), her fracture probability would be reduced to 13.6%.

The validity of the adjustment to FRAX has been explored in a meta-analysis of 14 cohorts (excluding Manitoba), together incorporating 17,809 men and women (59% women) ranging in age from 40 to 90 years (mean age 72 years) [42]. FRAX and TBS both had predictive ability for both major osteoporotic fractures and hip fracture, partly independently of each other. The predictive ability of FRAX was expressed as the gradient of risk (GR; hazard ratio per 1 SD change in risk variable in direction of increased risk). Overall, the GR of TBS for major osteoporotic fracture was 1.44 (95% CI: 1.35–1.53) when adjusted for age and time since baseline and was similar in men and women ($p > 0.10$). When additionally adjusted for FRAX 10-year probability of major osteoporotic fracture, TBS remained a significant, independent predictor for fracture (GR 1.32, 95% CI: 1.24–1.41). The adjustment of FRAX probability for TBS resulted in a small increase in the GR (1.76, 95% CI: 1.65–1.87 vs. 1.70, 95% CI: 1.60–1.81) suggesting that TBS would have clinical utility, for example, in the reclassification of those close to intervention thresholds.

Is TBS responsive to treatment?

Several relatively small studies have examined treatment-induced changes in TBS [49,50,53,56–59] (Table 3). Of these, four were studies of osteoporosis treatment in postmenopausal women, one in both

Table 3
Treatment related changes in lumbar TBS.

First author	Year published	Location	Study design	Subjects	TBS change/yr	Versus baseline	Versus other Rx	BMD change/yr	Versus baseline	Versus other Rx
Krieg [50]	2013	Canada	Case-control	534 PM women on standard OP Rx	0.20%	p < 0.001	p < 0.001	1.90%	p < 0.002	p < 0.001
				1150 untreated	−0.30%	p < 0.001		−0.40%	p < 0.001	
Popp [53]	2013	Switzerland	RCT	54 PM women on zoledronate	0.50%	p < 0.001	p < 0.001	3.20%	p < 0.001	p < 0.001
				53 on placebo	−0.20%	p < 0.001		0.50%	p < 0.001	
Senn [59]	2014	Switzerland	Open-label	65 PM women on teriparatide	2.20%	p < 0.001	p < 0.001	3.80%	p < 0.001	p < 0.001
				122 PM women on ibandronate	0.20%	p = 0.86		1.50%	p < 0.001	
Kalder [49]	2014	Germany	RCT substudy	17 PM women on tamoxifen	1.70%	p < 0.06	p < 0.001	1.00%	NS	p < 0.001
				19 PM women on exemestane	−1.20%	p < 0.05		−2.70%	p < 0.002	
Petranova [58]	2014	Bulgaria	Observational	36 PM women on denosumab	1.70%	p < 0.05	NA			
				No controls						
Kalder [57]	2015	Germany	RCT	34 women with breast cancer on zoledronate	2.41%	p < 0.006	p = 0.003	2.17%	p < 0.001	p < 0.005
				36 women with breast cancer on placebo	−0.52%			−5.02%		
Di Gregorio [56]	2015	Spain	Open-label	67 untreated	−1.55%	p < 0.05	NA	0.25%	NS	
				87 men and women on calcium and vitamin D	0.65%	NS	NA	0.80%	NS	NA
				88 men and women on alendronate	0.70%	p < 0.05	NA	2.05%	p < 0.001	NA
				36 men on testosterone	0.90%	NS	NA	2.20%	p < 0.001	NA
				39 men and women on risedronate	0.70%	NS	NA	2.40%	p < 0.001	NA
				43 men and women on denosumab	1.40%	p < 0.01	NA	4.40%	p < 0.001	NA
				30 men and women on teriparatide	1.80%	p < 0.01		4.40%	p < 0.001	NA

PM = postmenopausal; NA = not applicable.

men and women with osteoporosis, and two in the management of breast cancer. One of these compared a specific oestrogen-receptor modifier (tamoxifen) and an aromatase inhibitor (exemestane) in breast cancer patients.

In Manitoba, 534 post-menopausal women treated with a bisphosphonate (86%), raloxifene (10%) or calcitonin (4%), and having more than 75% compliance, were compared with 1150 untreated women [50]. The mean duration of follow-up was 3.7 years. Women in the treated group were older (mean age 66.1 vs. 62.2 years), had lower scores for aBMD and lumbar TBS, and had a higher prevalence of prior major osteoporotic fracture (15.4 vs. 10.4%) at baseline, consistent with indications for treatment. Over the course of follow-up, spine aBMD and TBS increased in treated women by 1.9% and 0.2%, respectively, whereas in untreated women aBMD and TBS decreased by 0.4% and 0.3% (all statistically significant changes versus baseline). Changes in aBMD and TBS from baseline were only weakly correlated ($r = 0.20$), consistent with the notion that the two indices represent partly independent measures of bone structure.

In the Swiss Horizon trial, 54 post-menopausal women treated with zoledronic acid were compared with 53 on placebo over 36 months of follow-up [52,53]. The only clinically-meaningful difference in baseline characteristics was a lower lumbar spine T-score in treated women (−2.9 vs. −2.1). Over three years, lumbar spine aBMD and TBS rose significantly, by 9.6% and 1.4%, respectively, in those on active treatment. The spine aBMD T-score also rose by 1.4% in those on placebo, perhaps due to degenerative disease, whilst TBS declined by 0.5%. The first statistically-significant change from baseline for aBMD was recorded at 6 months, whereas for TBS was at 24 months. Changes in the aBMD and TBS were only weakly correlated ($r = 0.20$) [52,53].

In a two-year open-label study [59] comparing teriparatide ($n = 65$) and ibandronate ($n = 122$), in which the only clinically-significant inter-group difference in baseline characteristics was a higher prevalence of past vertebral fractures in those on teriparatide (90.5 vs. 44.3%), patients on teriparatide had 7.6 and 4.3% increases in lumbar aBMD and TBS (both $p < 0.001$ vs. baseline), whilst only aBMD increased significantly in those on ibandronate (2.9% vs. 0.03%; $p < 0.001$ and 0.086, respectively). In this study, there was no significant correlation between changes in aBMD and TBS ($r^2 = 0.01$). In this report,

responsivity was also assessed as the proportion of patients achieving the least significant change in TBS and aBMD. For both treatment modalities lumbar spine BMD was more sensitive than TBS.

In an analysis of a small subset of treated female breast cancer patients in the Tamoxifen Exemestane Adjuvant Multicentre (TEAM) trial ($n = 19$ on exemestane, 17 on tamoxifen) [49], spine aBMD and TBS increased by 1.9 and 3.3% in those on tamoxifen versus a 5.3 and 2.3% decrease in patients given exemestane. The disparate responses to the two drugs are consistent with prior research demonstrating decreased fracture risk in postmenopausal breast cancer patients treated with tamoxifen and the increased risk observed in those on exemestane [60–62]. Changes in TBS and aBMD were again only weakly correlated ($r = 0.25$) [49].

A double-blind, randomized, placebo-controlled trial evaluated the effect of 8 cycles of adjuvant treatment with zoledronic acid over a 24-month period (4 mg i.v. once every 3 months) compared to placebo on aBMD and TBS in premenopausal women with oestrogen receptor and/or progesterone receptor positive breast cancer [57]. Treatment induced increases in both aBMD and TBS which were somewhat greater in percentage terms in the case of aBMD.

A cohort of 390 patients was analysed to evaluate the effect of different treatments (testosterone, risedronate, alendronate, denosumab, teriparatide) on aBMD and TBS [56]. After 24 months, a significant increase in TBS was observed only in the alendronate (+1.4%), denosumab (+2.8%) and teriparatide (+3.6%) groups, whereas aBMD increases in all treated groups. TBS was preserved on calcium and vitamin D, and decreased in the group without any treatment consistent with an epidemiological study showing that lower intakes of milk were associated with lower values for aBMD and TBS [63].

Longitudinal assessment of change represents a number of challenges both for bone densitometry and the evaluation of TBS. The precision of the measurement is critical, both within and between instruments [64]. However, sensitivity to measure changes is also affected by responsiveness, the ratio of responsiveness to precision being an indicator of sensitivity to measure (treatment-induced) changes [65]. At present, treatment induced changes in TBS have not been consistently assessed in this way. Nor has any possible contribution of TBS to fracture efficacy been explored, given that the change in aBMD with treatment

does not appear to fully explain the magnitude of fracture risk reduction [66,67].

Taken together, these intervention studies suggest that TBS tends to increase with treatments which increase aBMD, but that the magnitude of change of TBS is less marked than that of aBMD. It is currently unclear whether TBS might usefully contribute to the monitoring of treatment effects. However, these studies do indicate that TBS is potentially amenable to change as a result of pharmacological therapy. Whether this change is predictive of alterations in risk of future fracture, however, is currently not known.

Does TBS have a role in secondary osteoporosis?

Although in many cases, osteoporosis is idiopathic, there are a number of specific causes of bone fragility that result in “secondary osteoporosis”. Evidence is emerging that TBS might provide useful information with regards to bone health in several clinical contexts [68].

Clinical and subclinical hypercortisolism

Glucocorticoid induced osteoporosis (GIO) is common. Patients on long-term glucocorticoids fracture at a higher BMD than postmenopausal women, potentially implying a deleterious effect on bone structure. These effects may be more pronounced at lumbar spine; thus TBS may be well positioned to detect GIO-associated changes in bone structure.

In a cross-sectional study [65], women with systemic sclerosis were matched with 138 women with rheumatoid arthritis and 227 controls [69]. Multivariate analysis showed that a low TBS was independently associated with daily glucocorticoid dose. In a comparison of 34 patients with adrenal tumour and subclinical hypercortisolism [70], 68 patients with a tumour but no subclinical hypercortisolism, and 70 matched controls, both aBMD and TBS were lower in those with subclinical hypercortisolism. In addition, baseline lumbar TBS predicted incident fractures over a mean 40 months of follow-up, independent of patient age, BMI and lumbar aBMD [70].

Type-2 diabetes

There is evidence that patients with type-2 diabetes fracture at higher aBMD than postmenopausal women [71]. As such, there is a need for improved approaches to estimate fracture risk in such individuals. Recently, three studies have demonstrated that, although aBMD tends to be higher in type-2 diabetics than non-diabetics, the reverse is true of TBS. In the first paper, a cross-sectional case-control study by Dhaliwal et al., 57 women with type-2 diabetes were compared with 43 women without. TBS was lower and aBMD higher among diabetics ($p = 0.001$ and 0.01 , respectively). Moreover, TBS was lower ($p = 0.01$) and aBMD was no different in those diabetics with poor glycemic control compared to those with good glycemic control (an A1c above, versus below, 7.5%) [72]. The authors speculated that “abnormal architecture may help explain the paradox of increased fractures at higher aBMD” in these patients, although other mechanisms, such as glycosylation of collagen cross-links, key to the basic nanostructure of bone, have also been postulated [71]. Leslie et al. [73] retrospectively analysed data from 29,407 Canadian women age 50 years or over, who had undergone a baseline DXA examination, comparing 2356 diagnosed diabetics with the remainder of the cohort. After adjustment for clinical risk factors, diabetic women were found to be more likely to be in the lowest third of lumbar TBS, but less likely to be in the lowest thirds of lumbar, femoral neck or total hip aBMD. Both TBS and measures of aBMD were predictive of incident fracture. Finally, in the Ansong cohort, which included 1229 men and 1529 postmenopausal women (325 men and 370 women with type-2 diabetes) [74], lumbar spine TBS was lower in men and women with diabetes than those without, whereas lumbar spine BMD was higher in men and women

with diabetes. Interestingly, TBS was negatively correlated with HbA1c, fasting plasma glucose and fasting insulin. Based on the above studies, TBS may aid fracture risk assessment in patients with type-2 diabetes.

Primary hyperparathyroidism

Several studies provide support for use of lumbar TBS in the management of asymptomatic primary hyperparathyroidism [64,75–77]. Romagnoli et al. [77] noted lower TBS in 73 post-menopausal women with primary hyperparathyroidism (29 of them having a documented vertebral fracture) than in 74 age-matched controls. In a study that included both cross-sectional and longitudinal components, Eller-Vainicher et al. [76] compared 92 patients with primary hyperparathyroidism (74 of them post-menopausal females and the remainder males over age 50 years) and 98 controls with other conditions, consecutively recruited from clinic. Again, TBS was lower in patients with primary hyperparathyroidism than in controls, and was statistically significantly associated with vertebral fracture, even after adjustment for age, gender, BMI and lumbar spine aBMD (adjusted OR = 1.4; 95% CI: 1.1–1.9). In the longitudinal phase of the study, 20 primary hyperparathyroidism patients who underwent a parathyroidectomy to achieve ‘cure’ were compared at 24 months follow-up with 10 patients treated conservatively. In the surgery group, the mean TBS z-score increased by 1.20 ($p < 0.01$), whilst TBS non-significantly declined in the ten conservatively-treated counterparts [76].

Osteoarthritis

One limitation of spinal aBMD assessment is that the presence of overlying calcifications due to degenerative change can erroneously elevate the resulting measurement. Recent studies have suggested that TBS may be less affected by such artefactual influences [78]. Both points were demonstrated in a recent study by Kolta et al. [79] in which 1254 post-menopausal women in the European OPUS cohort, mean age 67 years, underwent TBS measurement at baseline and again at a mean of six years later. The severity of spinal osteoarthritis was graded according to Kellgren and Lawrence [80]. The investigators found that TBS was no different in women with, versus without, osteoarthritic changes in the spine at baseline, whilst lumbar spine aBMD measurements averaged 5.7% higher in the former group ($p < 0.003$). Over a mean six years of follow-up, TBS declined by 3.3% ($p < 0.001$), independent of the severity of spinal osteoarthritis. Conversely, though femoral neck and total hip aBMD also decreased over time and were not affected by osteoarthritis grade, there was no net decrease in lumbar aBMD, which was also associated with the grade of spinal arthritis [81].

Currently-published data support a role for the TBS in the assessment of fragility fracture risk in patients with a variety of secondary causes of osteoporosis, including subclinical hypercortisolism, type-2 diabetes, and parathyroid disease. Clearly, prospective cohort studies will be needed to confirm any added utility of TBS in these specific situations over and above its general predictive ability for fracture.

Conclusions

In recent years, there has been increasing interest in the use of TBS, a surrogate of bone microarchitecture, for risk stratification in osteoporosis. The present assessment of the existing literature indicates that low lumbar spine TBS is associated with both a history of fracture and the incidence of new fracture. The effect is independent of aBMD and of sufficient magnitude to enhance risk stratification with aBMD. The effect is also partly independent of FRAX with likely greatest utility for those individuals who lie close to an intervention threshold. TBS increases with treatment for osteoporosis, but the magnitude of this change is smaller than that with bone mineral density; the relationship between change in TBS and magnitude of fracture risk reduction remains to be elucidated. There have been a number of smaller investigations, which

have suggested that TBS may play a role in specific causes of increased fracture risk, such as glucocorticoid excess, hyperparathyroidism and type-2 diabetes.

Disclosures

NH has received consultancy, lecture fees and honoraria from the Alliance for Better Bone Health, AMGEN, MSD, Eli Lilly, Servier, Shire, Consilient Healthcare, Internis Pharma and Firefly Pharma.

CCG is a member of the Medimaps advisory board and has received consultancy fees from Medimaps.

RR has received speaker or advisory board fees from Amgen, MSD, GSK, Servier, Danone and Takeda.

NB has received research support from Amgen, Lilly, Merck, Opko Ireland, and GE Healthcare Lunar. He also declares consultancies/ advisory boards for Amgen, Lilly and Merck.

EM declares ad hoc consultancies/ speaking honoraria and/or research funding from Amgen, Bayer, General Electric, GSK, Hologic, Lilly, Merck Research Labs, Novartis, Novo Nordisk, Nycomed, Ono, Pfizer, ProStrakan, Roche, Sanofi-Aventis, Servier, Tethys, UBS and Warner-Chilcott.

MLB declares consultancies and grants from Alexion, Abiogen, Amgen, Bruno Farmaceutici, Eli Lilly, MSD, NPS, Shire, SPA and Servier.

CC has received consultancy, lecture fees and honoraria from AMGEN, GSK, Alliance for Better Bone Health, MSD, Eli Lilly, Pfizer, Novartis, Servier, Merck, Medtronic and Roche.

DLK has received consultancy, lecture fees, honoraria, and/or research grants from Amgen, Merck, Eli Lilly, GSK, Astalis, and Astra Zeneca. He holds shares in Medimaps.

OL has received speaker and/or advisory board fees and/or research grants from Amgen, Eli Lilly and Takeda.

AL has no interests to disclose.

BMC has received lecture fees from Achè, EMS, GSK, Grupogen-AC Farmacêutica, Hypermarcas, Lilly, Novartis, MSD, Sanofi, União Química, and Zodiac. No stock holding or shares to declare.

J-YR declares consulting fees or paid advisory boards from Servier, Novartis, Negma, Lilly, Wyeth, Amgen, GlaxoSmithKline, Roche, Merckle, Nycomed, NPS, Theramex, UCB; lecture fees from Merck Sharp and Dohme, Lilly, Rottapharm, IBSA, Genevrier, Novartis, Servier, Roche, GlaxoSmithKline, Teijin, Teva, Ebewee Pharma, Zodiac, Analis, Theramex, Nycomed, Novo-Nordisk, Nolver; Grant Support from Bristol Myers Squibb, Merck Sharp & Dohme, Rottapharm, Teva, Lilly, Novartis, Roche, GlaxoSmithKline, Amgen, and Servier.

JAK declares ad hoc consultancies/speaking honoraria and/or research funding from Amgen, D3A, Lilly, Medimaps, Merck Research Labs, Roche, Servier.

Acknowledgments

We are grateful for the assistance of Dr. Kevin P White (medical writer).

References

- [1] Anonymous. Consensus development conference. Diagnosis, prophylaxis and treatment of osteoporosis. *Am J Med* 1993;94:646–50.
- [2] World Health Organization. Assessment of fracture risk and its application to screening for postmenopausal osteoporosis: technical report series 843. Geneva: World Health Organization; 1994.
- [3] Kanis J, McCloskey EV, Johansson H, Oden A, Melton LJ, Khaltaev N. A reference standard for the description of osteoporosis. *Bone* 2008;42:467–75.
- [4] Kanis JA, on behalf of the World Health Organization Scientific Group. Assessment of osteoporosis at the primary health-care level. Technical Report. UK: World Health Organization Collaborating Centre for Metabolic Bone Diseases, University of Sheffield; 2007.
- [5] Miller PD, Siris ES, Barrett-Connor E, Faulkner KG, Wehren LE, Abbott TA, et al. Prediction of fracture risk in postmenopausal white women with peripheral bone densitometry: evidence from the National Osteoporosis Risk Assessment. *J Bone Miner Res* 2002;17:2222–30.
- [6] Albrand G, Munoz F, Sornay-Rendu E, Delmas PD. Independent predictors of all osteoporosis-related fractures in healthy postmenopausal women: the OFELY study. *Bone* 2003;32:78–85.
- [7] Hui SL, Slemenda CW, Johnson Jr CC. Age and bone mass as predictors of fracture in a prospective study. *J Clin Invest* 1988;81:1804–9.
- [8] Kanis JA, Johansson H, Oden A, Johnell O, De Laet C, Eisman JA, et al. A family history of fracture and fracture risk: a meta-analysis. *Bone* 2004;35:1029–37.
- [9] Kanis JA, Johnell O, De Laet C, Johansson H, Oden A, Delmas P, et al. A meta-analysis of previous fracture and subsequent fracture risk. *Bone* 2004;35:375–82.
- [10] Díez-Pérez A, González-Macías J, Marín F, Abizanda M, Alvarez R, Gimeno A, et al. Prediction of absolute risk of non-spinal fractures using clinical risk factors and heel quantitative ultrasound. *Osteoporos Int* 2007;18:629–39.
- [11] Bala Y, Zebaze R, Ghasem-Zadeh A, Atkinson EJ, Luliano S, Peterson JM, et al. Cortical porosity identifies women with osteopenia at increased risk for forearm fractures. *J Bone Miner Res* 2014;29:1356–62.
- [12] Faulkner KG, Cummings SR, Black D, Palermo L, Gluer CC, Genant HK. Simple measurement of femoral geometry predicts hip fracture: the study of osteoporotic fractures. *J Bone Miner Res* 1993;8:1211–7.
- [13] Gluer CC. Quantitative ultrasound – it is time to focus research efforts. *Bone* 2007;40:9–13.
- [14] Guerri-Fernandez RC, Noguez X, Quesada Gomez JM, Torres Del PE, Puig L, Garcia-Giralt N, et al. Microindentation for in vivo measurement of bone tissue material properties in atypical femoral fracture patients and controls. *J Bone Miner Res* 2013;28:162–8.
- [15] Keaveny TM, Kopperdahl DL, Melton III LJ, Hoffmann PF, Amin S, Riggs BL, et al. Age-dependence of femoral strength in white women and men. *J Bone Miner Res* 2010;25:994–1001.
- [16] Randall C, Bridges D, Guerri R, Noguez X, Puig L, Torres E, et al. Applications of a new handheld reference point indentation instrument measuring bone material strength. *J Med Devices* 2013;7:410051–6.
- [17] Silva BC, Leslie WD, Resch H, Lamy O, Lesnyak O, Binkley N, et al. Trabecular bone score: a noninvasive analytical method based upon the DXA image. *J Bone Miner Res* 2014;29:518–30.
- [18] Yang L, Peel N, Clowes JA, McCloskey EV, Eastell R. Use of DXA-based structural engineering models of the proximal femur to discriminate hip fracture. *J Bone Miner Res* 2009;24:33–42.
- [19] Cohen A, Dempster DW, Müller R, Guo XE, Nickolas TL, Liu XS, et al. Assessment of trabecular and cortical architecture and mechanical competence of bone by high-resolution peripheral computed tomography: comparison with transiliac bone biopsy. *Osteoporos Int* 2015;21:263–73.
- [20] Graeff C, Marin F, Petto H, Kayser O, Reisinger A, Peña J, et al. High resolution quantitative computed tomography-based assessment of trabecular microstructure and strength estimates by finite-element analysis of the spine, but not DXA, reflects vertebral fracture status in men with glucocorticoid-induced osteoporosis. *Bone* 2013;52:568–77.
- [21] Díez-Pérez A, Güerri R, Noguez X, Cáceres E, Peña MJ, Mellibovsky L, et al. Microindentation for in vivo measurement of bone tissue mechanical properties in humans. *J Bone Miner Res* 2010;25:1877–85.
- [22] Wang Y, Sanyal A, Cawthon PM, Palermo L, Jekir M, Christensen J, et al. Prediction of new clinical vertebral fractures in elderly men using finite element analysis of CT scans. *J Bone Miner Res* 2012;27:808–16.
- [23] Kopperdahl D, Aspelund T, Hoffman PF, Sigurdsson S, Siggeirsdottir K, Harris TB, et al. Assessment of incident spine and hip fractures in women and men using finite element analysis of CT scans. *J Bone Miner Res* 2014;29:570–80.
- [24] Silva BC, Leslie WD, Resch H, Lamy O, Lesnyak O, Binkley N, et al. Trabecular bone score: a noninvasive analytical method based upon the DXA image. *J Bone Miner Res* 2014;29:518–30.
- [25] Hans D, Goertzen AL, Krieg M-A, Leslie WD. Bone micro-architecture assessed by TBS predicts osteoporotic fractures independent of bone density: the Manitoba study. *J Bone Miner Res* 2011;26:2762–9.
- [26] Winzenrieth R, Michelet F, Hans D. Three-dimensional (3D) microarchitecture correlations with 2D projection image gray-level variations assessed by trabecular bone score using high-resolution computed tomographic acquisitions: effects of resolution and noise. *J Clin Densitom* 2013;16:287–96.
- [27] Simonelli C, Leib E, Mossman N, Winzenrieth R, Hans D, McClung M. Creation of an age-adjusted, dual-energy X-ray absorptiometry-derived trabecular bone score curve for the lumbar spine in non-Hispanic US White women. *J Clin Densitom* 2014;17:314–9.
- [28] Del Rio LM, Winzenrieth R, Cormier C, Di GS. Is bone microarchitecture status of the lumbar spine assessed by TBS related to femoral neck fracture? A Spanish case-control study. *Osteoporos Int* 2013;24:991–8.
- [29] Krueger D, Fidler E, Libber J, Aubry-Rozier B, Hans D, Binkley N. Spine trabecular bone score subsequent to bone mineral density improves fracture discrimination in women. *J Clin Densitom* 2014;17:60–5.
- [30] Lamy O, Krieg MA, Stoll D, Aubry-Rozier B, Metzger M, Hans D. The OsteoLaus Cohort Study: bone mineral density, microarchitecture score and vertebral fracture assessment extracted from a single DXA device in combination with clinical risk factors improve significantly the identification of women at high risk of fracture. *Osteologie* 2012;21:77–82.
- [31] Leib E, Winzenrieth R, Aubry-Rozier B, Hans D. Vertebral microarchitecture and fragility fracture in men: a TBS study. *Bone* 2014;62:51–5.
- [32] Leib E, Winzenrieth R, Lamy O, Hans D. Comparing bone microarchitecture by trabecular bone score (TBS) in Caucasian American women with and without osteoporotic fractures. *Calcif Tissue Int* 2014;95:201–8.
- [33] Nassar K, Paternotte S, Kolta S, Fechtenbaum J, Roux C, Briot K. Added value of trabecular bone score over bone mineral density for identification of vertebral

- fractures in patients with areal bone mineral density in the non-osteoporotic range. *Osteoporos Int* 2014;25:243–9.
- [34] Pothuaud L, Barthe N, Krieg MA, Mehsen N, Carceller P, Hans D. Evaluation of the potential use of trabecular bone score to complement bone mineral density in the diagnosis of osteoporosis: a preliminary spine BMD-matched, case-control study. *J Clin Densitom* 2009;12:170–6.
- [35] Rabier B, Winzenrieth R, Héraud A, Grand-Lenoir C, Pothuaud L, Hans D. A multicentre, retrospective case-control study assessing the role of trabecular bone score (TBS) in a cohort of menopausal Caucasian women with low bone mineral density (BMD): analysing the odds of vertebral fracture. *Bone* 2008;46:176–81.
- [36] Touvier J, Winzenrieth R, Johansson H, Roux JP, Chaintreuil J, Toumi H, et al. Fracture discrimination by combined bone mineral density (BMD) and microarchitectural texture analysis. *Calcif Tissue Int* 2015;96:274–83.
- [37] Vasic J, Petranova T, Povoroznyuk V, Barbu CG, Karadzic M, Gozkovic F, et al. Evaluating spine micro-architectural texture (via TBS) discriminates major osteoporotic fractures from controls both as well as and independent of site matched BMD: the Eastern European TBS study. *Bone Miner Metab* 2014;32:556–62.
- [38] Winzenrieth R, Dufour R, Pothuaud L, Hans D. A retrospective case-control study assessing the role of trabecular bone score in postmenopausal Caucasian women with osteopenia: analyzing the odds of vertebral fracture. *Calcif Tissue Int* 2010;86:104–9.
- [39] Boutroy S, Hans D, Sornay-Rendu E, Vilayphiou N, Winzenrieth R, Chapurlat R. Trabecular bone score improves fracture risk prediction in non-osteoporotic women: the OFELY study. *Osteoporos Int* 2013;24:77–85.
- [40] Briot K, Paternotte S, Kolta S, Eastell R, Reid DM, Felsenberg D, et al. Added value of trabecular bone score to bone mineral density for prediction of osteoporotic fractures in postmenopausal women: the OPUS study. *Bone* 2013;57:232–6.
- [41] Iki M, Tamaki J, Kadowaki E, Sato Y, Dongmei N, Winzenrieth R, et al. Trabecular bone score (TBS) predicts vertebral fractures in Japanese women over 10 years independently of bone density and prevalent vertebral deformity: the Japanese Population-based Osteoporosis (JPOS) cohort study. *J Bone Miner Res* 2014;29:399–407.
- [42] Kanis JA, Oden A, Harvey NC, Leslie WD, Hans D, Johansson H, et al. A meta-analysis of trabecular bone score in fracture risk prediction and its interaction with FRAX. *Osteoporos Int* 2015;26.
- [43] Leslie WD, Johansson H, Kanis JA, Lamy O, Oden A, McCloskey EV, et al. Lumbar spine texture enhances 10-year fracture probability assessment. *Osteoporos Int* 2014;25:2271–7.
- [44] Leslie WD, Lix LM. Comparison between various fracture risk assessment tools. *Osteoporos Int* 2014;25:1–21.
- [45] Hans D, Goertzen AL, Krieg MA, Leslie WD. Bone microarchitecture assessed by TBS predicts osteoporotic fractures independent of bone density: the Manitoba study. *J Bone Miner Res* 2011;26:2762–9.
- [46] Leslie WD, Caetano PA, MacWilliam LR, Finlayson GS. Construction and validation of a population-based bone densitometry database. *J Clin Densitom* 2005;8:25–30.
- [47] Leslie WD, Anderson WA, Metge CJ, Manness LJ. Clinical risk factors for fracture in postmenopausal Canadian women: a population-based prevalence study. *Bone* 2007;40:991–6.
- [48] Hans D, Barthe N, Boutroy S, Pothuaud L, Winzenrieth R, Krieg MA. Correlations between trabecular bone score, measured using anteroposterior dual-energy X-ray absorptiometry acquisition, and 3-dimensional parameters of bone microarchitecture: an experimental study on human cadaver vertebrae. *J Clin Densitom* 2011;14:302–12.
- [49] Kalder M, Hans D, Kyveritakis I, Lamy O, Bauer M, Hadji P. Effects of exemestane and tamoxifen treatment on bone texture analysis assessed by TBS in comparison with bone mineral density assessed by DXA in women with breast cancer. *J Clin Densitom* 2014;17:66–71.
- [50] Krieg MA, Aubry-Rozier B, Hans D, Leslie WD. Effects of anti-resorptive agents on trabecular bone score (TBS) in older women. *Osteoporos Int* 2013;24:1073–8.
- [51] Popp A, Meer S, Krieg M-A, Perrelet R, Hans D, Lippuner K. Bone mineral density (BMD) and vertebral trabecular bone score (TBS) for the identification of elderly women at high risk for fracture: the SEMOF cohort study. *Eur Spine J* 2015 [In press].
- [52] Muschitz C, Kocijian R, Pahr D, Patsch JM, Amrein K, Misof BM, Kaider A, Resch H, Pietschmann P. Ibandronate increases sclerostin levels and bone strength in male patients with idiopathic osteoporosis. *Calcif Tissue Int* 2015;96:477–89.
- [53] Popp AW, Guler S, Lamy O, Senn C, Buffat H, Perrelet R, et al. Effects of zoledronate versus placebo on spine bone mineral density and microarchitecture assessed by the trabecular bone score in postmenopausal women with osteoporosis: a three-year study. *J Bone Miner Res* 2013;28:449–54.
- [54] Leslie WD, Aubry-Rozier B, Lix LM, Morin SN, Majumdar SR, Hans D. Spine bone texture assessed by trabecular bone score (TBS) predicts osteoporotic fractures in men: the Manitoba Bone Density Program. *Bone* 2014;67:10–4.
- [55] McCloskey EV, Oden A, Harvey NC, Leslie WD, Hans D, Johansson H, et al. Adjusting fracture probability by trabecular bone score. *Calcif Tissue Int* 2015;96:500–9.
- [56] Di Gregorio S, Del Rio L, Rodriguez-Tolra J, Bonel E, García M, Winzenrieth R. Comparison between different bone treatments on areal bone mineral density (aBMD) and bone microarchitectural texture as assessed by the trabecular bone score (TBS). *Bone* 2015;75:138–43.
- [57] Kalder M, Kyveritakis I, Albert US, Baier-Ebert M, Hadji P. Effects of zoledronic acid versus placebo on bone mineral density and bone texture analysis assessed by the trabecular bone score in premenopausal women with breast cancer treatment-induced bone loss: results of the ProBONE II substudy. *Osteoporos Int* 2015;26:353–60.
- [58] Petranova T, Sheytanov I, Monov S, Rashkov R, Kinov P. Efficacy of denosumab for control of bone mineral density and microarchitecture in postmenopausal women with osteoporosis: a one-year study. *Osteoporos Int* 2015;26:3977–81.
- [59] Senn C, Gunther B, Popp AW, Perrelet R, Hans D, Lippuner K. Comparative effects of teriparatide and ibandronate on spine bone mineral density (BMD) and microarchitecture (TBS) in postmenopausal women with osteoporosis: a 2-year open-label study. *Osteoporos Int* 2014;25:1945–51.
- [60] Zidan J, Keidar Z, Basher W, Israel O. Effects of tamoxifen on bone mineral density and metabolism in postmenopausal women with early-stage breast cancer. *Med Oncol* 2004;21:117–21.
- [61] Barni S, Lissoni P, Tancini G, Ardizzoia A, Cazzaniga M. Effects of one-year adjuvant treatment with tamoxifen on bone mineral density in postmenopausal breast cancer women. *Tumori* 1996;82:65–7.
- [62] Love RR, Mazess RB, Barden HS, Epstein S, Newcomb PA, Jordan VC, et al. Effects of tamoxifen on bone mineral density in postmenopausal women with breast cancer. *N Engl J Med* 1992;326:852–6.
- [63] Sato Y, Iki M, Fujita Y, Tamaki J, Kouda K, Yura A, et al. Greater milk intake is associated with lower bone turnover, higher bone density, and higher bone microarchitecture index in a population of elderly Japanese men with relatively low dietary calcium intake: Fujiwara-kyo Osteoporosis Risk in Men (FORMEN) Study. *Osteoporos Int* 2015;26:1585–94.
- [64] Silva BC, Bilezikian JP. Trabecular bone score: perspectives of an imaging technology coming of age. *Arq Bras Endocrinol Metabol* 2014;58:493–503.
- [65] Gluer CC. Monitoring skeletal changes by radiological techniques. *J Bone Miner Res* 1999;14:1952–62.
- [66] Watts NB, Geusens P, Barton IP, Felsenberg D. Relationship between changes in BMD and nonvertebral fracture incidence associated with risendronate: reduction in risk of nonvertebral fracture is not related to change in BMD. *J Bone Miner Res* 2015;20:2097–104.
- [67] Watts NB, Cooper C, Lindsay R, Eastell R, Manhart MD, Barton IP, et al. 2004 Relationship between changes in bone mineral density and vertebral fracture risk associated with risendronate: greater increases in bone mineral density do not relate to greater decreases in fracture risk. *J Clin Densitom* 2004;7:255–61.
- [68] Ulivieri FM, Silva BC, Sardanelli F, Hans D, Bilezikian JP, Caudarella R. Utility of the trabecular bone score (TBS) in secondary osteoporosis. *Endocrine* 2014;47:435–48.
- [69] Koumakis E, Winzenrieth R, Guignat L, Cormier C. Cushing disease: gain in bone mineral density and also bone texture assessed by trabecular bone score after cure of Cushing disease. *J Bone Miner Res* 2014;29(Suppl. 1):S349.
- [70] Eller-Vainicher C, Morelli V, Ulivieri FM, Palmieri S, Zhukouskaya VV, Cairoli E, et al. Bone quality, as measured by trabecular bone score in patients with adrenal incidentalomas with and without subclinical hypercortisolism. *J Bone Miner Res* 2012;27:2223–30.
- [71] Saito M, Kida Y, Kato S, Marumo K. Diabetes, collagen, and bone quality. *Curr Osteoporos Rep* 2014;12:181–8.
- [72] Dhaliwal R, Cibula D, Ghosh C, Weinstock RS, Moses AM. Bone quality assessment in type 2 diabetes mellitus. *Osteoporos Int* 2014;25:1969–73.
- [73] Leslie WD, Aubry-Rozier B, Lamy O, Hans D. TBS (trabecular bone score) and diabetes-related fracture risk. *J Clin Endocrinol Metab* 2013;98:602–9.
- [74] Kim JH, Choi HJ, Ku EJ, Kim KM, Kim SW, Cho NH, et al. Trabecular bone score as an indicator for skeletal deterioration in diabetes. *J Clin Endocrinol Metab* 2015;100:475–82.
- [75] Bilezikian JP, Brandi ML, Eastell R, Silverberg SJ, Udelman R, Marcocci C, et al. Guidelines for the management of asymptomatic primary hyperparathyroidism: summary statement from the Fourth International Workshop. *J Clin Endocrinol Metab* 2014;99:3561–9.
- [76] Eller-Vainicher C, Filopanti M, Palmieri S, Ulivieri FM, Morelli V, Zhukouskaya VV, et al. Bone quality, as measured by trabecular bone score, in patients with primary hyperparathyroidism. *Eur J Endocrinol* 2013;169:155–62.
- [77] Romagnoli E, Cipriani C, Nofroni I, Castro C, Angelozzi M, Scarpello A, et al. “Trabecular bone score” (TBS): an indirect measure of bone micro-architecture in postmenopausal patients with primary hyperparathyroidism. *Bone* 2013;53:154–9.
- [78] Dufour R, Winzenrieth R, Heraud A, Hans D, Mehsen N. Generation and validation of a normative, age-specific reference curve for lumbar spine trabecular bone score (TBS) in French women. *Osteoporos Int* 2013;24:2837–46.
- [79] Kolta S, Briot K, Fechtenbaum J, Paternotte S, Ambrecht G, Felsenberg D, et al. TBS result is not affected by lumbar spine osteoarthritis. *Osteoporos Int* 2014;25:1759–64.
- [80] Yoshida K, Barr RJ, Galea-Soler S, Aspden RM, Reid DM, Gregory JS. Reproducibility and diagnostic accuracy of Kellgren-Lawrence grading for osteoarthritis using radiographs and dual-energy X-ray absorptiometry images. *J Clin Densitom* Oct 8 2014. <http://dx.doi.org/10.1016/j.jocd.2014.08.003> [pii: S1094-6950(14)00206-6, Epub ahead of print].
- [81] Breban S, Briot K, Kolta S, Paternotte S, Ghazi M, Fechtenbaum J, et al. Identification of rheumatoid arthritis patients with vertebral fractures using bone mineral density and trabecular bone score. *J Clin Densitom* 2012;15:260–6.